What is Claimed

1. A method of treating bulimia nervosa in a mammal, comprising administering to the mammal an effective amount of a compound of the formula:

$$R_5$$
 R_7
 R_7
 R_7


5

10

15

20

in which A is a moiety of the formula



wherein

the dotted line represents optional unsaturation;

R1 is hydrogen or alkyl of 1 to 6 carbon atoms;

R₂ is alkyl of 1 to 6 carbon atoms;

R4 is hydrogen, alkyl of 1 to 6 carbon atoms, formyl, or alkanol of 2 to 7 carbon atoms;

R5 and R6 are, independently, hydrogen, hydroxyl, alkyl of 1 to 6 carbon atoms, alkoxy of 1 to 6 carbon atoms, alkanoyloxy of 2 to 7 carbon atoms, cyano, nitro, alkylmercapto of 1 to 6 carbon atoms, amino, alkylamino of 1 to 6 carbon atoms, dialkylamino in which each alkyl group is of 1 to 6 carbon atoms, alkanamido of 2 to 7 carbon atoms, halo, trifluoromethyl, or taken together, methylene dioxy;

R7 is hydrogen or alkyl of 1 to 6 carbon atoms; and

n is 0, 1, 2, 3, or 4;

or a pharmaceutically acceptable salt thereof.


- 2. The method of claim 1 wherein the mammal is a human.
- 3. The method of claim 1 wherein the composition is administered orally.

)

4. A method of treating bulimia nervosa in a mammal, comprising administering to the mammal a pharmaceutical composition comprising an effective amount of a compound of the formula:

$$R_5$$
 R_7
 R_7
 R_7

in which A is a moiety of the formula



10

15

20

25

wherein

the dotted line represents optional unsaturation, and

R₁ is hydrogen or alkyl of 1 to 6 carbon atoms;

R₂ is alkyl of 1 to 6 carbon atoms;

R4 is hydrogen, alkyl of 1 to 6 carbon atoms, formyl, or alkanol of 2 to 7 carbon atoms:

R5 and R6 are, independently, hydrogen, hydroxyl, alkyl of 1 to 6 carbon atoms, alkoxy of 1 to 6 carbon atoms, alkanoyloxy of 2 to 7 carbon atoms, cyano, nitro, alkylmercapto of 1 to 6 carbon atoms, amino, alkylamino of 1 to 6 carbon atoms, dialkylamino in which each alkyl group is of 1 to 6 carbon atoms, alkanamido of 2 to 7 carbon atoms, halo, trifluoromethyl, or taken together, methylene dioxy;

R7 is hydrogen or alkyl of 1 to 6 carbon atoms; and

n is 0, 1, 2, 3, or 4;

or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier or excipient.

5. The method of claim 4 wherein the compound is:

$$R_5$$
 R_7
 R_7
 R_7

5 in which A is a moiety of the formula

wherein

10

15

20

the dotted line represents optional unsaturation, and

R₁ is hydrogen or alkyl of 1 to 3 carbon atoms;

R₂ is alkyl of 1 to 3 carbon atoms;

R₅ is hydrogen, hydroxyl, alkoxy of 1 to 3 carbon atoms, chloro, bromo, trifluoromethyl or alkyl of 1 to 3 carbon atoms;

R₆ is alkyl of 1 to 3 carbon atoms, alkoxy of 1 to 3 carbon atoms, chloro, bromo, trifluoromethyl or alkanoyloxy of 2 to 3 carbon atoms;

R7 is hydrogen or alkyl of 1 to 3 carbon atoms;

or a pharmaceutically acceptable salt thereof.

6. The method of claim 1 wherein R_5 and R_6 are both in the meta positions or one of R_5 or R_6 is in the para position and n is 2.

7. The method of claim 1 wherein the compound is 1-[(2-dimethylamino)-1-(4-methoxyphenyl)ethyl]cyclohexanol or a pharmaceutically acceptable salt thereof.

10

15

- 8. The method of claim 1 wherein the compound is 1-[2-(dimethylamino)-1-(4-hydroxyphenyl)ethyl]cyclohexanol or a pharmaceutically acceptable salt thereof.
- 5 9. The method of claim 1 wherein the effective amount comprises a daily dose of between about 50 mg/day and about 375 mg/day.
 - 10. The method of claim 1 wherein the effective amount comprises a daily dose of between about 75 mg/day and about 200 mg/day.
 - 11. The method of claim 1 wherein the mammal is a human.
 - 12. The method of claim 1 wherein the pharmaceutical composition is administered orally.
 - 13. The method of claim 1 wherein the pharmaceutical composition is in a unit dosage form which is a tablet.
- 14. The method of claim 1 wherein the pharmaceutical composition is in a20 unit dosage form which is a capsule.